

AD-A041 168

NAVAL POSTGRADUATE SCHOOL MONTEREY CALIF
EFFICIENT COMPUTATION AND LONG RANGE OPTIMIZATION APPLICATIONS --ETC(U)
MAY 77 K T MARSHALL
NPS55-77-23

F/6 12/2

UNCLASSIFIED

NL

1 OF 1
AD
A041168



ADA 041 168

NPS55-77-23

NAVAL POSTGRADUATE SCHOOL
Monterey, California



EFFICIENT COMPUTATION AND LONG RANGE
OPTIMIZATION APPLICATIONS USING
A TWO-CHARACTERISTIC MARKOV-TYPE
MANPOWER FLOW MODEL


by

Kneale T. Marshall

May 1977

Approved for public release; distribution unlimited.

Prepared for:
Headquarters Marine Corps, Code MPI20
Washington, D.C. 20380



AD No. _____
DDC FILE COPY


NAVAL POSTGRADUATE SCHOOL
Monterey, California

Rear Admiral Isham Linder
Superintendent

Jack R. Borsting
Provost


This work was supported in part by the Manpower Planning Division (MPI20) of the Marine Corps through the Navy Personnel Research and Development Lab., San Diego.


Reproduction of all or part of this report is authorized.


Kneale T. Marshall
Professor
Department of Operations Research

Reviewed by:

Released by:


Michael G. Sovereign, Chairman
Department of Operations Research


Robert R. Fossum
Dean of Research

Unclassified

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER NPS55-77-23	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) Efficient Computation and Long Range Optimization Applications Using a Two-Characteristic Markov-Type Manpower Flow Model		5. TYPE OF REPORT & PERIOD COVERED Technical <i>report</i>
7. AUTHOR(s) Kneale T. Marshall		6. PERFORMING ORG. REPORT NUMBER
9. PERFORMING ORGANIZATION NAME AND ADDRESS Naval Postgraduate School Monterey, California 93940		8. CONTRACT OR GRANT NUMBER(s)
11. CONTROLLING OFFICE NAME AND ADDRESS Naval Postgraduate School Monterey, California 93940		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS N6822177WR70052
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		12. REPORT DATE May 1977
		13. NUMBER OF PAGES 18
		15. SECURITY CLASS. (of this report) Unclassified
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report) Approved for public release; distribution unlimited		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Manpower; Personnel Optimization; Markov		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) In the past, the author has compared and contrasted Markov and longitudinal manpower flow models. The Markov model requires relatively little data and has been widely analyzed, (see [2] and [3]). The longitudinal model incorporates more realistic personnel flows, but requires extensive data which is not always available. In [4] Hayne and Marshall analyzed a two-characteristic Markov model which can be viewed as a hybrid of the Markov and longitudinal models. The purpose of this paper is to show how efficient computational		

DD FORM 1 JAN 73 1473

EDITION OF 1 NOV 65 IS OBSOLETE
S/N 0102-014-6601

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

251 450

next page

679

20. methods can be used with the two-characteristic model by exploiting the special structure of its underlying matrix. These methods make possible the efficient use of this basic flow model in optimization models similar to those described in Chapter 5 of [3].

ACCESSION for	
NTIS	White Section <input checked="" type="checkbox"/>
DTIC	Soft Section <input type="checkbox"/>
UNANNOUNCED	<input type="checkbox"/>
JUSTIFICATION.....	
BY.....	
DISTRIBUTION/AVAILABILITY CODES	
Dist.	AVAIL. AND/OR SPECIAL
A	

DDC
 RECORDED
 JUL 5 1977
 RECEIVED
 - D -

EFFICIENT COMPUTATION AND LONG RANGE OPTIMIZATION APPLICATIONS
USING A TWO-CHARACTERISTIC MARKOV-TYPE
MANPOWER FLOW MODEL

by

Kneale T. Marshall
Naval Postgraduate School
Monterey, CA 93940

ABSTRACT

In [1] the author has compared and contrasted Markov and longitudinal manpower flow models. The Markov model requires relatively little data and has been widely analyzed (see [2] and [3]). The longitudinal model incorporates more realistic personnel flows, but requires extensive data which is not always available. In [4] Hayne and Marshall analyze a two-characteristic Markov model which can be viewed as a hybrid of the Markov and longitudinal models. The purpose of this paper is to show how efficient computational methods can be used with the two-characteristic model by exploiting the special structure of its underlying matrix. These methods make possible the efficient use of this basic flow model in optimization models similar to those described in Chapter 5 of [3].

D D C
RECEIVED
JUL 5 1977
REGISTERED
D

TABLE OF CONTENTS

	Page
1. Introduction	1
2. Model Formulation	2
3. The Structure of $\tilde{\ell}(\alpha)$ and $\tilde{P}(\alpha)$	5
4. Computation of $\tilde{\ell}(\alpha)$ and $\tilde{P}(\alpha)$	11
5. Numerical Example	15
References	18

1. Introduction

In a previous paper [1] the author has compared and contrasted cross-sectional and longitudinal manpower flow models. The cross-sectional, or Markov, model requires relatively little data and has been widely analyzed (see, for example, [2] and [3]). The longitudinal model incorporates more realistic personnel flows, but requires extensive data which is not always available. In [4] Hayne and Marshall analyze a two-characteristic cross-sectional model which can be viewed as a hybrid of the Markov and longitudinal models. In Chapter 5 of [3] Grinold and the author present some long-range optimization models based primarily on the longitudinal model.

The purpose of this paper is to show how efficient computational methods can be used with the two-characteristic model by exploiting the special structure of its underlying matrix. These methods make possible the efficient use of this basic flow model in optimization models similar to those described in [3]. This paper explores this application in detail when the two characteristics of the state are grade and time-in-grade. Approximate solutions are found to infinite horizon linear programs using methods similar to those in [3].

This paper relies heavily on the notation and results in [3] and [4]. The reader is referred there for details. Section 2 contains a formulation of the optimization model and its approximation (see [3], Chapter 5,

pages 186-209). Section 3 contains results on the structure of the flow matrices, their generating function for the (Grade, Time in Grade) model, and the legacies and their generating function. Section 4 describes efficient methods of computation for these generating functions. Section 5 gives a simple numeric example which the reader may wish to follow simultaneously with the theory in Sections 3 and 4. The example has been kept small and simple because of space limitations and for ease of exposition.

2. Model Formulation

It is assumed that manpower joins a system on one of K chains, and at some discrete time t is counted in one of n classes if it is still in the system. Let $P(u)$ be an $n \times K$ matrix with element $p_{ij}(u)$ equal to the fraction of manpower that enters on chain j which is in class i , u periods after entering the system. The matrices $P(0), P(1), \dots$ describe the flow through the system.

Let $g(u)$ be a K -element column vector of flows into the system on each chain at time $u = 1, 2, \dots$; let $\ell(u)$ be an n -element column vector of legacies in each class at time $u = 1, 2, \dots$ of manpower which enter up to and including time $t = 0$, which is taken to be the current time. If $s(t)$ is an n -element vector of stocks in each class at time $t = 1, 2, \dots$, then

$$(1) \quad s(t) = \sum_{u=1}^t P(t-u) g(u) + \ell(t) .$$

In [3] Grinold and Marshall used this chain-flow model as the basis of a linear optimization model (see Chapter 5). Let α be a discount factor, a an n -element vector of one-period costs on stocks, b a K -element of one-period costs on new hires, ρ the constant size of the system, A a constraint matrix on stocks and B a constraint matrix on flows. Finally let e be a vector with all elements equal to 1. Consider the infinite horizon linear program (LP)

$$\begin{array}{ll} \text{Minimize} & \sum_{t=1}^{\infty} \alpha^t [as(t) + bg(t)] \\ \text{Pl.} & \\ \text{Subject to:} & \left. \begin{array}{l} es(t) = \rho \\ As(t) \geq 0 \\ Bg(t) \geq 0 \\ g(t) \geq 0 \end{array} \right\} \quad t = 1, 2, \dots \end{array}$$

It is shown in [3] that the solution to a K -variable single period problem can be used to generate solutions to Pl which are usually optimal and always good approximations. Let

$$\tilde{P}(\alpha) = \sum_{u=0}^{\infty} \alpha^u P(u), \quad \tilde{\ell}(\alpha) = \sum_{u=1}^{\infty} \alpha^u \ell(u) \text{ and } c = a\tilde{P}(\alpha) + b.$$

Then by using (1) to eliminate $s(t)$, and multiplying the t -th period constraints in Pl by α^t and summing, we obtain

$$\begin{aligned}
\text{P2.} \quad & \text{Minimize } cg \\
& \text{Subject to: } e\tilde{P}(\alpha)g = [\alpha\rho/(1-\alpha)] - e\tilde{\ell}(\alpha) \\
& \quad \tilde{A}P(\alpha)g \geq -\tilde{A}\tilde{\ell}(\alpha) \\
& \quad Bg \geq 0 \\
& \quad g \geq 0
\end{aligned}$$

This LP has only K variables g . Let g^* be the optimal solution to P2. In [3], Chapter 5, it is shown how scalars $\gamma(1), \gamma(2), \dots$ can easily be found successively such that, if we let $g^*(t) = \gamma(t)g^*$, then these $g^*(t)$ are often optimal in P1 (they are optimal if A is vacuous).

The purpose of this paper is to analyze the structure of $\tilde{P}(\alpha)$ and $\tilde{\ell}(\alpha)$ in P2 when

- (i) The system can be partitioned into a hierarchical structure of grades
- (ii) A two-characteristic cross-sectional flow model is used with a state described by the 2-tuple (Grade, Time in Grade). This is called the (G, TIG)-model.
- (iii) Entering a chain corresponds to entering a grade, necessarily with TIG equal to 1.
- (iv) A class corresponds to a grade.

We use the results of Hayne and Marshall in [4] for the (G, TIG) model. For example, in a simple model of a university faculty

the grades might be Assistant Professor, Associate Professor and Full Professor. Let us assume that the maximum number of periods (years) a person can spend in these grades is 6, 30, and 30. The number of states in the (G,TIG) model is 66, so that a cross-sectional flow matrix, say Q , (66×66) would have 4356 elements, most of which would have value zero. One of the matrices $P(u)$ in (1) would be 3×3 with only 9 elements, since chains map onto grades with TIG equal to 1 and stocks are also measured only in grades. Any given manpower policy will imply certain element values for Q . In this paper we show how to find $P(u)$ and $\ell(u)$, and hence $\tilde{P}(\alpha)$ and $\tilde{\ell}(\alpha)$, from Q in an efficient manner. Thus policy changes can be reflected in the chain flow model through Q , where the effects of such changes on $P(u)$ could not be determined directly. A simple numerical example is given in Section 5 following the theory. The reader may wish to follow the example concurrently with the theory.

3. The Structure of $\tilde{\ell}(\alpha)$ and $\tilde{P}(\alpha)$

The number of grades in the system is $n (=K)$. Let the maximum time in grade j be $u(j)$, and let $\ell = \sum_{j=1}^n u(j)$. Then ℓ is the number of states in the system. Let Q be the one-period flow matrix for the (G,TIG) model. The u -period flows are given by Q^u . We now relate the $\ell \times \ell$ matrix Q^u to the $n \times n$ matrix $P(u)$.

Let Γ be an $n \times \ell$ matrix where row j has

- (i) the first $\sum_{i=1}^{j-1} u(i)$ elements equal to 0,
- (ii) the next $u(j)$ elements equal to 1,
- (iii) the remaining elements equal to 0.

Let Φ be an $\ell \times n$ matrix where column j has

- (i) the $(u(j-1)+1)$ -th element equal to 1 (element 1 when $j = 1$)
- (ii) all other elements equal to zero.

Then

$$(2) \quad P(u) = \Gamma Q^u \Phi, \quad u = 0, 1, 2, \dots$$

Recall that $\tilde{P}(\alpha) = \sum_{u=0}^{\infty} \alpha^u P(u)$. Using (2), if we let

$$(3) \quad N(\alpha) = (I - \alpha Q)^{-1}$$

then

$$(4) \quad \tilde{P}(\alpha) = \Gamma N(\alpha) \Phi.$$

Recall also that $\tilde{\ell}(\alpha) = \sum_{u=1}^{\infty} \alpha^u \ell(u)$.

Let σ be the ℓ -vector of stocks of manpower in each state at time $t = 0$. Then

$$\ell(u) = \Gamma Q^u \sigma, \quad u = 1, 2, \dots$$

and therefore,

$$(5) \quad \tilde{\ell}(\alpha) = \alpha \Gamma (N(\alpha) - I) \sigma.$$

Notice from (4) and (5) that to determine $\tilde{P}(\alpha)$ and $\tilde{\ell}(\alpha)$ we need to compute the $n \times \ell$ matrix $\Gamma N(\alpha)$. But first we look at the structure of $N(\alpha)$, the inverse of the large sparse matrix $(I - \alpha Q)$.

In [4] Hayne and Marshall show that for the (G,TIG) model the cross-sectional flow matrix Q has the structure

$$(6) \quad Q = \begin{bmatrix} Q_1 & & & & & \\ P_1 & Q_2 & & & & \\ & \cdot & \cdot & & & \\ & & \cdot & \cdot & & \\ & & & \cdot & P_{n-1} & \cdot \\ & & & & Q_n & \end{bmatrix}$$

where Q_j is a $u(j) \times u(j)$ matrix and P_j is a $u(j+1) \times u(j)$ matrix. All other submatrices of Q have 0's as elements and are suppressed. In addition each Q_j contains zeros except for the lower diagonal,

$$(7) \quad Q_j = \begin{bmatrix} 0 & \dots\dots\dots & 0 \\ q_{j1} & 0 & & & & \\ 0 & q_{j2} & \cdot & & & \\ \cdot & \cdot & \cdot & \cdot & & \\ \cdot & & \cdot & \cdot & \cdot & \\ \cdot & & & \cdot & \cdot & \cdot \\ 0 & \dots\dots\dots & 0 & q_{j,u(j)-1} & 0 \end{bmatrix},$$

$$j = 1, 2, \dots, n.$$

Also each P_j contains zeros except for the top row,

$$(8) \quad P_j = \begin{bmatrix} p_{j1} & p_{j2} & \cdots & p_{j,u(j)} \\ 0 & & & 0 \\ \vdots & & & \vdots \\ 0 & \text{.....} & & 0 \end{bmatrix}, \quad j = 1, 2, \dots, n-1.$$

The inverse of $(I - \alpha Q)$ can be written

$$(9) \quad N(\alpha) = \begin{bmatrix} N_{11}(\alpha) & & & & \\ N_{21}(\alpha) & \cdot & & & \\ \cdot & \cdot & \cdot & & \\ \cdot & \cdot & \cdot & \cdot & \\ \cdot & & \cdot & \cdot & \cdot \\ N_{n1}(\alpha) & & N_{n,n-1}(\alpha) & & N_{nn}(\alpha) \end{bmatrix}$$

where

$$N_{jj}(\alpha) = (I - \alpha Q_j)^{-1}$$

and

$$N_{ji} = \alpha^{j-i} N_{jj}(\alpha) (P_{j-1} N_{j-1,j-1}(\alpha)) \cdots (P_i N_{ii}(\alpha))$$

for $j > i$. Thus $N(\alpha)$ is completely determined by the inverses $\{N_{jj}(\alpha), j = 1, 2, \dots, n\}$.

Now let

$$(10) \quad n_{ki}(j) = q_{j,k-1} \cdot q_{j,k-2} \cdots \cdot q_{j,i} \quad \text{for } k > i,$$

the product of $(k-i)$ nonzero elements of Q_j . Then

$$(11) \quad N_{jj}(\alpha) = \begin{bmatrix} 1 & & & & & \\ \alpha n_{21}(j) & 1 & & & & \\ \alpha^2 n_{31}(j) & \alpha n_{32}(j) & 1 & & & \\ \cdot & \cdot & \cdot & \cdot & \cdot & \\ \cdot & \cdot & \cdot & \cdot & \cdot & \\ \cdot & \cdot & \cdot & \cdot & \cdot & \\ \alpha^{n-1} n_{u(j),1}(j) & & & & \alpha n_{u(j),u(j-1)}(j) & 1 \end{bmatrix}$$

$$i = 1, 2, \dots, n,$$

Thus all the elements of $N_{jj}(\alpha)$ are determined from the partial products in (10), and all the elements of $N(\alpha)$ in (9) are determined by multiplication of these with the vectors forming the top rows of the P_j in (8). However, the matrix $N(\alpha)$ in (9) need never be explicitly determined in order to find $\tilde{P}(\alpha)$ and $\tilde{l}(\alpha)$. We make the following observations.

- $$(12) \quad \left\{ \begin{array}{l} (i) \quad P_j N_{jj}(\alpha) \text{ is a matrix with nonzero elements only} \\ \text{in the top row (i.e., it has the same structure} \\ \text{as (8)).} \\ (ii) \quad \text{If } \Pi \text{ is any matrix with the same structure as} \\ \text{(8) with as many rows as } N_{jj}(\alpha) \text{ has columns,} \\ \text{then } N_{jj}(\alpha) \Pi \text{ requires only the first column of} \\ N_{jj}(\alpha) \end{array} \right.$$

Now let

$$r = \text{Max}\{u(j), j = 1, 2, \dots, n\},$$

and let $W(\alpha)$ be an $n \times r$ matrix with the j -th row equal to the nonzero elements of Q_j multiplied by α , preceded by a 1, and 0's added to the right as necessary. Thus

$$(13) \quad W(\alpha) = \begin{bmatrix} 1 & \alpha q_{11} & \alpha q_{12} & \cdots & \alpha q_{1, u(1)-1} & \cdots & 0 & \cdots & 0 \\ 1 & \alpha q_{21} & \alpha q_{22} & \cdots & \alpha q_{2, u(2)-1} & 0 & \cdots & 0 \\ \vdots & & & & & & & & \\ 1 & \alpha q_{n1} & \alpha q_{n2} & \cdots & \alpha q_{n, u(n)-1} & \cdots & 0 & \cdots & 0 \end{bmatrix}$$

where it has been assumed that $u(1) < u(2) < u(n) = r$ for clarity. Similarly, let $V(\alpha)$ be an $n \times r$ matrix with the j -th row equal to the nonzero elements of P_j multiplied by α , with 0's added to the right as necessary, for $j \leq n-1$, and row n a row of 0's. Thus

$$(14) \quad V(\alpha) = \begin{bmatrix} \alpha p_{11} & \cdots & \alpha p_{1, u(1)} & \cdots & 0 & \cdots & 0 \\ \alpha p_{21} & \cdots & \alpha p_{2, u(2)} & 0 & \cdots & 0 \\ \vdots & & & & & & \\ \alpha p_{n1} & \cdots & \alpha p_{n, u(n)} & \cdots & 0 & \cdots & 0 \\ 0 & \cdots & 0 & \cdots & 0 & \cdots & 0 \end{bmatrix}$$

Finally let S be an $n \times r$ matrix of stocks at time zero, where element s_{ij} gives the stocks in grade i with time in grade equal to j . Thus S is a matrix representation of the vector σ with 0's added where necessary. Let $T(j) = \sum_{i=1}^j u(i)$. Then

$$(15) \quad S = \begin{bmatrix} \sigma_1 & \sigma_2 & \dots & \sigma_{T(1)} & 0 & \dots & 0 \\ \sigma_{T(1)+1} & \sigma_{T(1)+2} & \dots & \sigma_{T(2)} & 0 & \dots & 0 \\ \vdots & & & & & & \vdots \\ \sigma_{T(n-1)+1} & \sigma_{T(n-1)+2} & \dots & \dots & \dots & \dots & \sigma_{T(n)} \end{bmatrix}$$

All the information required to determine $\tilde{\ell}(\alpha)$ and $\tilde{P}(\alpha)$ is contained in $W(\alpha)$, $V(\alpha)$, and S . Both can be found using simple row operations on these compact matrices. In the 3 grade, 66 state example given earlier V , W and S are each 3×30 and contain a total of 270 elements. Storage of the Q matrix to find $(I - \alpha Q)^{-1}$ directly would require 4356 elements, and direct inversion would necessitate inverting a 66 dimensional very sparse matrix. Total storage for both Q and σ would be 4422 elements.

4. Computation of $\tilde{\ell}(\alpha)$ and $\tilde{P}(\alpha)$.

Let v and w be any two n -vectors. We define a vector-valued function $F(v, w)$ on these such that

$$(16) \quad F(v, w) = [(v_1 + w_2 v_2 + \dots + w_2 w_3 \dots w_n v_n), (v_2 + w_3 v_3 + \dots + w_3 w_4 \dots w_n v_n), \dots, (v_{n-1} + w_n v_n), v_n] .$$

We can extend this function as follows. If V and W are matrices of the same size, then $F(V,W)$ is a matrix X with the same dimensions as V and W , and if v and w are the j -th rows of V and W , respectively, then (16) gives the j -th row of X . The function F is the key to the efficient calculation of both $\tilde{l}(\alpha)$ and $\tilde{P}(\alpha)$, since by using it we can efficiently compute $\Gamma N(\alpha)$.

Let U be an $n \times r$ matrix such that element (i,j) is 1 if element (i,j) of $W(\alpha)$ is positive, and is 0 otherwise, where $W(\alpha)$ is given by (13). Also let

$$v_j = e N_{jj}(\alpha), \quad j = 1, 2, \dots, n,$$

the column sums of $N_{jj}(\alpha)$ in (11). Thus v_j is a $u(j)$ -vector. Now let \bar{v}_j be v_j followed by $r-u(j)$ zero's. Then

$$(17) \quad F(U, W(\alpha)) = \begin{bmatrix} \bar{v}_1 \\ \bar{v}_2 \\ \vdots \\ \bar{v}_n \end{bmatrix},$$

an $n \times r$ matrix.

Now let m_j be the top row of the matrix $P_j N_{jj}(\alpha)$, $j = 1, 2, \dots, n-1$, a $u(j)$ -vector, and let \bar{m}_j be m_j followed by $r-u(j)$ zero's. Then

$$(18) \quad F(V(\alpha), W(\alpha)) = \begin{bmatrix} \bar{m}_1 \\ \bar{m}_2 \\ \vdots \\ \bar{m}_{n-1} \\ 0 \end{bmatrix},$$

an $n \times r$ matrix. Equations (17) and (18) contain all the required data for $\Gamma N(\alpha)$. Let v^1 be the first element of any vector v . Then using the results in (11) and (12),

$$(19) \quad \Gamma N(\alpha) = \begin{bmatrix} v_1 & & & & & \\ v_2^1 m_1 & v_2 & & & & \\ v_3^1 m_2 m_1 & v_3^1 m_2 & v_3 & & & \\ \vdots & & & \ddots & & \\ v_n^1 m_{n-1}^1 \cdots m_2^1 m_1 & \cdots \cdots \cdots & v_n & & & \end{bmatrix},$$

an $n \times l$ matrix.

Consider now the determination of $\tilde{P}(\alpha) = \Gamma N(\alpha) \Phi$. From (19) and the definition of Φ we see that

$$(20) \quad \tilde{P}(\alpha) = \begin{bmatrix} v_1^1 & & & & \\ v_2^{1m_1} & v_2^1 & & & \\ v_3^{1m_2m_1} & v_3^{1m_2} & v_3^1 & & \\ \vdots & & & \ddots & \\ v_n^{1m_{n-1} \dots m_2 m_1} & \dots & \dots & \dots & v_n^1 \end{bmatrix},$$

an $n \times n$ matrix. This can be constructed from the first columns of (17) and (18).

To determine $\tilde{\ell}(\alpha)$ we require $\Gamma N(\alpha)$ and σ . From (19) it is easy to see that it can be constructed from (17), (18), and (15).

We now illustrate efficient *APL* functions to compute the various matrices. The reader should consult [5] for details of *APL* programming.

Let $M1$ and $M2$ be any two matrices of the same dimension. Then the *APL* function

$$(21) \quad \begin{aligned} &V \leftarrow M1 \ F \ M2; X \\ [1] \quad &R \leftarrow (\neq \times \backslash M2 + 0 = M2) \times \phi + \backslash \phi M1 \times X + 0 = X \leftarrow \times \backslash M2 \\ &V \end{aligned}$$

calculates $F(M1, M1)$ as defined by (16). Let V , W and S be *APL* variables for the matrices $V(\alpha)$, $W(\alpha)$ and S in (14), (13), and (15) respectively. Then (17) is given by $(W > 0) \ F \ W$, and (18) by $V \ F \ W$. If we let these be X and Y respectively,

and let N be the number of grades and P be $\tilde{P}(\alpha)$, then

$$(22) \quad P \leftarrow ((1N) \circ . \geq 1N) \times ((N, N) \rho X[;1]) \times \backslash (x \div 1, Z), [1] \rho (N, N-1) \rho Z \leftarrow 1 + Y[;1]$$

Finally, if A is the APL variable for α , and L the variable for $\tilde{l}(\alpha)$, then from (5), and the arguments above

$$(23) \quad L \leftarrow A \times ((+/X \times S) + (P \times ((1N) \circ . > 1N) \times (N, N) \rho \div Z, 1) + . \times + / Y \times S) \div + / S.$$

5. Numerical Example

Assume we have 3 grades and that the maximum time in each grade is 3, 4 and 5 periods. Thus $n = 3$, $u(1) = 3$, $u(2) = 4$, $u(3) = 5$, $r = 5$. The matrix Q in (6) is 12×12 and contains the following 5 non-zero submatrices (see (7) and (8))

$$Q_1 = \begin{bmatrix} 0 & 0 & 0 \\ .9 & 0 & 0 \\ 0 & .8 & 0 \end{bmatrix}$$

$$P_1 = \begin{bmatrix} .05 & .1 & .8 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}$$

$$Q_2 = \begin{bmatrix} 0 & 0 & 0 & 0 \\ .95 & 0 & 0 & 0 \\ 0 & .9 & 0 & 0 \\ 0 & 0 & .85 & 0 \end{bmatrix}$$

$$P_2 = \begin{bmatrix} 0 & 0 & .1 & .7 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

$$Q_3 = \begin{bmatrix} 0 & 0 & 0 & 0 & 0 \\ .9 & 0 & 0 & 0 & 0 \\ 0 & .9 & 0 & 0 & 0 \\ 0 & 0 & .8 & 0 & 0 \\ 0 & 0 & 0 & .7 & 0 \end{bmatrix}$$

The starting stocks are given in the 12-vector

$$\sigma = (100, 73, 70, 82, 65, 63, 58, 59, 48, 30, 25, 20) .$$

Let $\alpha = 0.9$. Then from (13), (14) and (15) we have the 3×5 matrices

$$W(\alpha) = \begin{bmatrix} 1 & .81 & .72 & 0 & 0 \\ 1 & .855 & .81 & .765 & 0 \\ 1 & .81 & .81 & .72 & .63 \end{bmatrix} ,$$

$$V(\alpha) = \begin{bmatrix} .045 & .09 & .72 & 0 & 0 \\ 0 & 0 & .09 & .63 & 0 \\ 0 & 0 & 0 & 0 & 0 \end{bmatrix} ,$$

$$S = \begin{bmatrix} 100 & 73 & 70 & 0 & 0 \\ 82 & 65 & 63 & 58 & 0 \\ 59 & 48 & 30 & 25 & 20 \end{bmatrix} .$$

From (17) and (18), by using (21) we obtain

$$F(U, W(\alpha)) = \begin{bmatrix} 2.393 & 1.72 & 1 & 0 & 0 \\ 3.077 & 2.429 & 1.765 & 1 & 0 \\ 3.236 & 2.761 & 2.174 & 1.63 & 1 \end{bmatrix}$$

and

$$F(V(\alpha), W(\alpha)) = \begin{bmatrix} .538 & .608 & .72 & 0 & 0 \\ .398 & .463 & .472 & .63 & 0 \\ 0 & 0 & 0 & 0 & 0 \end{bmatrix}$$

Then using (22) and (23)

$$\tilde{P}(\alpha) = \begin{bmatrix} 2.393 & 0 & 0 \\ 1.655 & 3.077 & 0 \\ 0.689 & 1.282 & 3.236 \end{bmatrix}$$

and

$$\tilde{\ell}(\alpha) = (172.7 \quad 691.9 \quad 805.8) .$$

Once $\tilde{P}(\alpha)$ and $\tilde{l}(\alpha)$ have been determined they can be used in the optimization problem P2 to find optimum input flows each period. Suppose we wish to measure the effect of a change of policy. Let us assume for example that we want to test the effect of limiting the maximum time in grade 2 to 3 periods, and of extending the maximum time in grade 3 to 6 periods. Then $W(\alpha)$, $V(\alpha)$, and S each become 3×6 matrices. But (21) can be used with any sized matrices and so (22) and (23) readily give the new values of $\tilde{P}(\alpha)$ and $\tilde{l}(\alpha)$ for use in resolving P2. The computational speeds involved in these calculations, including solving P2 and determining $g^*(t) = \gamma(t)g$, are of the order 1 or 2 seconds for systems with about 10 grades and a maximum time in grade of about 30 periods. Thus interactive computer models can be designed so that the manpower planner can sit at a terminal and test alternate policies or design manpower systems.

REFERENCES

- [1] Marshall K. T., "A Comparison of Two Personnel Prediction Models," Operations Research 21, 810-822 (1973).
- [2] Bartholomew, D. J., Stochastic Models for Social Processes, 2nd edition, John Wiley, New York, 1973.
- [3] Grinold, R. C. and Marshall, K. T., Manpower Planning Models, Elsevier—North Holland, New York, 1977.
- [4] Hayne, W. J. and Marshall, K. T., "Two-Characteristic Markov-type Manpower Flow Models," report NPS55MT74071, 1974, Naval Postgraduate School, Monterey, California 93940 (to appear in Nav. Res. Log. Quart. (1977)).
- [5] Gilman, L. and Rose, A. J., APL: An Interactive Approach, John Wiley, New York, 1974.